



After Your Kidney Biopsy

1. You may remove the Band-Aid from the biopsy site the morning after your biopsy.
2. You may shower the morning after your biopsy.
3. Do not do strenuous exercise or activity for 5 days after the biopsy.
4. If you have pain, take Tylenol 325 mg: two tablets every 4 to 6 hours. If pain is not relieved, or if it persists for more than 48 hours, call the transplant center at the numbers above.
5. Call your doctor if you experience:
 - fever
 - chills
 - redness or tenderness at the biopsy site
 - drainage at the biopsy site
 - difficulty urinating or inability to urinate
 - bleeding from the site
 - blood in your urine
 - foul-smelling urine
 - abdominal pain
6. Other instructions:

Contacts (Reach the staff members below through the NIH page operator at **301-496-1211**.)

Your Doctor:

| |
|--|
| |
| |
| |



2004

This information is prepared specifically for patients participating in clinical research at the Warren Grant Magnuson Clinical Center at the National Institutes of Health and is not necessarily applicable to individuals who are patients elsewhere. If you have questions about the information presented here, talk to a member of your healthcare team.

Questions about the Clinical Center? OCCC@cc.nih.gov

Where applicable, brand names of commercial products are provided only as illustrative examples of acceptable products, and do not imply endorsement by NIH; nor does the fact that a particular brand name product is not identified imply that such product is unsatisfactory.

National Institutes of Health
Warren Grant Magnuson Clinical Center
Bethesda, Maryland 20892